

5. Данилов В. И., Недосекина Т.В. О влиянии разных режимов содержания степи на длительное сохранения степной растительности в условиях заповедника «Галичья гора» // Проблемы сохранения и восстановления степных ландшафтов : материалы Межрегиональных научных чтений. – Оренбург, 1999. – С. 48.
6. Дидух Я. П., Лысенко Г. Н. Экологические проблемы охраны степей Украины // Степи Евразии : проблемы сохранения и восстановления. – С.-Пб.–М. : Ин-т географ. РАН, 1993. – С. 65 – 77.
7. Жерихин В. В. Природа и история травяных биомов // Степи Евразии: проблемы сохранения и восстановления. – С.Пб.–М. : Институт географии РАН, 1993. – С. 29 – 49.
8. Захаренко А. В., Грамма В. Н. К вопросу об управлении экосистемами степи // Современные проблемы заповедного дела : тез. докл. обл. научн. конф. – Курск, 1985. – С. 33 – 35.
9. Краснитский А. М. Проблемы заповедного дела. – М.: Лесная промышленность, 1983. – 191 с.
10. Лысенко Г. Н. Влияние абсолютно заповедного режима на изменение гидротермических и эдафических факторов экотопов «Михайловской целины» (Украина) // Степи Евразии : сохранение природного разнообразия и мониторинг состояния экосистем. Материалы международного симпозиума. – Оренбург, 1997. – С. 53 – 54.
11. Лысенко Г. Н. В каком режиме сохранится луговая степь «Михайловской целины»? // Степной бюллетень. – 2005. – № 18. – С. 10 – 14.
12. Лысенко Г. Н. Луговая степь «Михайловская целина» (Украина): проблема выбора режимов заповедания // История заповедного дела : мат-лы междунар. научн. конфер. – Борисовка, 2005. – С. 133 – 136.
13. Лысенко Г. Н. Степные заповедники Украины как флористические инсультные геосистемы // Живые объекты в условиях антропогенного пресса: материалы X Международной научно-практической экологической конференции (г. Белгород, 15-18 сентября 2008 г.). – Белгород : ИПЦ «ПОЛИТЕРРА», 2008. – С. 122.
14. Национальна доповідь про стан навколишнього природного середовища в Україні у 2012 році. – К. : Міністерство екології та природних ресурсів України, Олді-плюс. – 2014. – 282 с.
15. Осичнюк В. В. Деякі особливості заповідного режиму у відділеннях Українського державного степового заповідника // Укр. ботан. журн. – 1979. – Т. 36, № 4. – С. 347 – 352.
16. Петрович О. З., Іваненко Є. І., Драпалюк А. М. Аналіз структури та територіальної репрезентативності Природно-заповідного фонду України // Збірник наук. праць ДНБС. – 2013. – Т. 135. – С. 7 – 16.
17. Разумовский С. М. Закономерности динамики биоценозов. – М. : Наука, 1981. – 231 с.
18. Семенова-Тян-Шанская А. М. Накопление и роль подстилки в травяных сообществах. – Л. : Наука, 1977. – 191 с.
19. Ткаченко В. С. Фітоценологічний моніторинг резерватних сукцесій в Українському степовому природному заповіднику. – К. : Фітосоціоцентр, 2004. – 184 с.
20. Ткаченко В. С., Лысенко Г. Н. Комплексные мероприятия по охране луговой степи «Михайловской целины» // Проблемы сохранения разнообразия природы степных и лесостепных регионов. Материалы российско-украинской науч. конф. посвящ. 60-летию Центрально-Черноземного заповедника (пос. Заповедный, Курск. обл., 22 – 27 мая 1995 г.). – М. : КМС Scientific Press Ltd., 1995. – С. 74 – 76.

УДК 632.118.3: 581.557.24

THE ROLE OF ARBUSCULAR MYCORRHIZAL SYMBIOSIS IN ¹³⁴CS UPTAKE BY CROP AND WILD PLANT SPECIES

Sergiy Dubchak

Ph.D. in biology, Associate Professor, State Ecological Academy of Postgraduate Education and Management, V. 03035, V.Lypkivsky str. 35, build. 2, sergiy.dubchak@yahoo.com

The role of arbuscular mycorrhizal fungus *Glomus intraradices* in ¹³⁴Cs isotope uptake by different plant species is studied. The impact of radiocaesium on mycorrhizal development and functioning of plant photosynthetic apparatus is considered. The possibility of mycorrhizal symbiosis application in phytoremediation of radioactively contaminated areas is analyzed. It is found that colonization of plants by arbuscular mycorrhizal fungus resulted to significant decrease of radiocaesium content in their aboveground parts, while it didn't have considerable impact on the radiocesium uptake by plant root system. *Keywords:* radiocaesium, radioactive contamination of environment, arbuscular mycorrhiza, arbuscular mycorrhizal fungi, plant photosynthetic apparatus, mycorrhizal colonization, phytoremediation.

Роль арбускулярного мікоризного симбіозу в накопиченні ¹³⁴Cs дикорослими та культурними видами рослин. Сергій Валерійович Дубчак. Досліджено роль арбускулярного мікоризного гриба *Glomus intraradices* у накопиченні ізотопу ¹³⁴Cs різними видами рослин. Розглянуто вплив радіоцезію на розвиток мікоризи та функціонування фотосинтетичного апарату рослин. Проаналізовано можливість застосування мікоризного симбіозу у фіторемерації радіаційно забруднених територій. Встановлено, що колонізація рослин арбускулярним мікоризним грибом призвела до суттєвого зменшення концентрації радіоцезію в їхній надземній частині й водночас не мала значного впливу на надходження радіонукліда до кореневої системи рослин. *Ключові слова:* радіоцезій, радіоактивне забруднення довкілля, арбускулярна мікориза, арбускулярні мікоризні гриби, фотосинтетичний апарат рослин, мікоризна колонізація, фіторемерація.

Роль арбускулярного мікоризного симбіозу в накопиченні ¹³⁴Cs дикорастущими та культурними видами рослин. Сергей Валериевич Дубчак. Исследована роль арбускулярного мікоризного гриба *Glomus intraradices* в накопиченні ізотопу ¹³⁴Cs різними видами рослин. Рассмотрено влияние радиоцезия на развитие микоризы и функционирование фотосинтетического аппарата растений. Проанализирована возможность применения микоризного симбіозу в фиторемерации радиоактивно загрязненных территорий. Установлено, что колонизация арбускулярным микоризным грибом привела к существенному уменьшению концентрации радиоцезия в их надземной части и одновременно не имела значительного влияния на поступления радионуклида в корневую систему растений. *Ключевые слова:* радиоцезий, радиоактивное загрязнение окружающей среды, арбускулярная микориза, арбускулярные микоризные грибы, фотосинтетический аппарат растений, микоризная колонизация, фиторемерация.

Statement of the problem. The radiocaesium isotopes have been introduced into the environment via various routes for last several decades. Altogether, roughly 1 EBq (10¹⁸ Bq) of long-lived ¹³⁷Cs was released to the Earth's biosphere in the XX – XXIth centuries that resulted to contamination of vast areas all over the

world. About 90 % of radiocaesium was originated from atmospheric nuclear testing, approximately 4 % was released by fuel reprocessing and nuclear fuel facilities and roughly 6 % – by Chernobyl and Fukushima accidents. Nowadays the ^{137}Cs absorption by plants and its accumulation, therefore, represents the main source of human exposure to this radionuclide. The principal route of radiocaesium entry into biological food chain in terrestrial ecosystems is the soil-to-plant pathway. This radionuclide is expected to remain in the rooting zone of plants for decades and respectively to be involved in biological migration chains. However, the mechanisms by which radiocaesium is taken up by plant roots are not completely understood.

Analysis of recent studies and publications. Recently the alternative strategies, orientated towards the use of plants and micro-organisms, separately or in combination, have been proposed for removing or immobilizing radiocaesium and other radionuclides in the soil [1,2]. Among these micro-organisms, mycorrhizal fungi received a particular attention. An estimated 90 % of terrestrial plants exist in a symbiotic association with soil fungi forming mycorrhizal associations. Among them, the obligate arbuscular mycorrhizal (AM) fungal symbionts are supposed to have a principal role [3]. These fungi are important participants in the Cs cycle in the upper layers of soils. They have strong impact on mobility of radiocaesium in the soil and result to unavailability of this radionuclide to the other components in ecosystems [4]. At the same time, it was demonstrated [5] that AM fungi can transform and immobilize radionuclides and correspondingly limit their toxicity and bioavailability to plants and spreading into the soils. Accordingly, plants

growing in contaminated soil could obtain benefit from their AM fungal symbiotic partners.

Nevertheless, the role of arbuscular mycorrhizal fungi on the acquisition of radiocaesium by plants remains poorly understood and controversial. The lack of clear results on the capacity of AM fungi to accumulate or transport Cs could be principally attributed to different and inadequate experimental systems used in previous studies. Furthermore, the various AM fungi and plants studied could also explain the controversial conclusions obtained, since AM fungi and plants have probably different capacity to accumulate and transport radiocaesium.

Objectives of research. The goals of this work were to identify the capacity of AM fungi to take up and transfer caesium isotopes to their hosts as well as to estimate the influence of arbuscular mycorrhiza on radiocaesium uptake by plants and impact of radiocaesium on development of AM fungal symbioses.

The main material of the study. Four plant species (*Plantago lanceolata*, *Medicago truncatula*, *Lolium perenne* and *Helianthus annuus*) capable to form efficient association with a broad range of AM fungi were selected for our study. The plants were cultivated in the presence or absence of AM fungus *Glomus intraradices* (strain BIO, obtained from BIORIZE, Dijon, France). The ^{134}Cs isotope (obtained from "POLATOM" Radioisotope Centre, Otwock-Świerk, Poland) in was added to sterilized substrata in pots in form of CsCl water solution. ^{134}Cs activity concentration was adjusted to 100 000 Bq per pot (77 000 Bq·kg⁻¹). The plants were grown in transparent Sun bags (SigmaTM Aldrich, Poznan, Poland) in a

growth chamber at 20 °C, with a photoperiod of 12 h light and 12 h darkness, at photosynthetic photon flux density $30 \pm 6 \mu\text{mol}\cdot(\text{s}\cdot\text{m}^2)^{-1}$ and harvested each three months.

The activity concentration of ^{134}Cs in roots and shoots of plants was determined using a gamma-spectrometer with semiconductor p-type coaxial high purity HP-Ge detector with a relative efficiency of 15 % and resolution of 2.5 keV at 1.33 MeV, shielded by 10 cm of lead with inner lining with 2 mm Cd and 18 mm Cu.

For the estimation of mycorrhizal colonization, the roots of plants were carefully washed with tap water, softened in 10% potassium hydroxide for 24 hours, washed in water again, acidified in 5% lactic acid in water for 12 – 24 h and stained with 0.01% aniline blue in lactic acid (to visualize AMF) for 24 h at room temperature. The root fragments were mounted and squashed on the slide covered with lactoglycerole. The parameters of AM colonization were assessed according to the method developed in [6] that assumes six levels of mycorrhizal colonization (from 0 to 5). The relative mycorrhizal root length (M%), intensity of colonization within individual mycorrhizal roots (m%), relative arbuscular richness (A%) and arbuscule richness in root fragments where the arbuscules were present (a%) were evaluated using Nikon Eclipse 800 light microscope equipped with Nomarski contrast and fluorescence.

The photosynthetic activity of plants was evaluated using a Plant Efficiency Analyzer fluorimeter (Hansatech Instruments, UK) estimating Chlorophyll *a* fluorescence transients of intact plant leaves. The Chl *a* fluorescence transients (OJIP transients) were induced by a

red light pulse (peak at 650 nm) of 600 W·m⁻² intensity provided by an array of three light-emitting diodes. The transients were recorded for 1 s with 12 bit resolution, starting 10 μs after the onset of illumination. Each transient was analyzed according to the OJIP-test based on the theory of energy fluxes in biomembranes [7]. The selected original data were processed by means of their utilization for the calculation of biophysical parameters by the JIP-test equation and the number of biophysical parameters were calculated. Among them, the most parameters are the performance indexes PI_{abs} (evaluated on the base of light absorption) and PI_{total} (total performance index). PI_{abs} and PI_{total} comply all basic biophysical parameters and represent the photosynthetic system vitality.

Part 1. ^{134}Cs uptake by plant species.

P. lanceolata inoculated with *G. intraradices* contained $66846 \pm 11029 \text{ Bq}\cdot\text{kg}^{-1}$ of ^{134}Cs in their shoots, that is considerably lower in comparison with the radionuclide activity concentration in nonmycorrhizal plant shoots ($87500 \pm 12333 \text{ Bq}\cdot\text{kg}^{-1}$). At the same time, ^{134}Cs activity concentration in roots of mycorrhizal and nonmycorrhizal *P. lanceolata* wasn't differed significantly, although the slightly higher radiocaesium concentration ($18.1 \pm 9.9 \%$) was found in roots of nonmycorrhizal plants (Fig. 1A). Due to the higher biomass of mycorrhizal *P. lanceolata*, the ^{134}Cs activity in roots and shoots of single mycorrhizal and nonmycorrhizal plant (Bq·plant⁻¹, dry weight) and correspondingly the total radiocaesium activity in a single plant

(i.e. shoots plus roots) weren't differed substantially ($p < 0.05$). The root/shoot ratios of ^{134}Cs activity concentration in *P. lanceolata* colonized with *G. intraradices* were slightly (about 11 %) higher as compared to those of nonmycorrhizal ones. As it's known [8], the higher root/shoot ratios of caesium content in plants indicates the reduced root to shoot translocation of this element, thus the tendency of the mycorrhiza to reduce radionuclide translocation from *P. lanceolata* roots to shoots was revealed.

The colonization of *M. truncatula* with *G. intraradices* also caused a significant reduction of radiocaesium uptake in plant shoots. Thus, ^{134}Cs activity concentration in aboveground part of mycorrhizal *M. truncatula* was $86888 \pm 20022 \text{ Bq}\cdot\text{kg}^{-1}$, whereas shoots of nonmycorrhizal plants contained $132100 \pm 15505 \text{ Bq}\cdot\text{kg}^{-1}$ of this radionuclide. At the same time, the mycorrhiza resulted to considerable ($18.8 \pm 5.6 \%$) increase of radiocaesium activity concentration in *M. truncatula* roots in comparison with that in nonmycorrhizal plants (see Fig. 2B). The distribution of ^{134}Cs activity between aboveground and underground parts of mycorrhizal and nonmycorrhizal *M. truncatula* was differed. Thus, the radionuclide activity concentration in roots of mycorrhizal *M. truncatula* was $45.5 \pm 14.7 \%$ lower than in their shoots. The opposite tendency was observed in case of nonmycorrhizal plants, where ^{134}Cs activity concentration in shoots was $24.2 \pm 8.9 \%$ higher than in roots (see Fig. 2B). No statistically significant differences were found between dry masses of mycorrhizal and nonmycorrhizal plants, although both roots and shoots of AM inoculated *M. truncatula* grown on ^{134}Cs spiked substrata had slightly higher weight (about 14 and

12 % correspondingly) as compared to those of nonmycorrhizal plants. The colonization of plants grown on radioactively contaminated substrata also led to moderate (about 10 %) increase of their shoot length. The evaluated ^{134}Cs activity in shoots of single mycorrhizal alfalfa was $9.3 \pm 0.6 \text{ Bq}$, whereas shoots of nonmycorrhizal *M. truncatula* contained significantly higher amount of radiocaesium ($12.3 \pm 0.6 \text{ Bq}$). On the contrary, the radionuclide activity in roots of mycorrhizal alfalfa was substantially higher ($3.1 \pm 0.1 \text{ Bq}$) as compared to that of nonmycorrhizal plants ($2.3 \pm 0.2 \text{ Bq}$). Consequently, mycorrhizal *M. truncatula* had significantly lower total activity of radiocaesium ($12.4 \pm 0.7 \text{ Bq}$) when compared to that of nonmycorrhizal plants ($14.6 \pm 0.8 \text{ Bq}$). The radionuclide translocation from underground to aboveground parts of plants was more intensive in case of nonmycorrhizal alfalfa. Their root/shoot ratio of ^{134}Cs activity concentration was 0.81 ± 0.28 being considerably lower in comparison with that of mycorrhizal *M. truncatula* (1.45 ± 0.41).

The harvested plants of *L. perenne* mycorrhizal with *G. intraradices* had more than two fold lower ^{134}Cs activity concentration both in their roots and shoots as compared to those of nonmycorrhizal plants. The ^{134}Cs distribution within *L. perenne* demonstrated that the radionuclide activity concentration in aboveground parts of both mycorrhizal and nonmycorrhizal ryegrass was about three times lower when compared to that of plant underground parts (see Fig. 1C). Dry weights of mycorrhizal and nonmycorrhizal *L. perenne* and their shoot length weren't differed considerably, however the biomass of plants colonized with *G. intraradices* and grown on substrata

spiked with ^{134}Cs was slightly (less than 10 %) higher when compared to that of nonmycorrhizal species. Hence, the colonization with *G. intraradices* resulted to considerable decrease of ^{134}Cs activity (Bq per plant) in shoots ($76.3 \pm 22.8 \%$) and roots ($53.3 \pm 15.8 \%$) of single mycorrhizal ryegrass as compared to that of nonmycorrhizal plants.

As opposed to plant species considered above, the colonization of *H. annuus* with *G. intraradices* resulted to significant increase of ^{134}Cs uptake by plants. Thus, the radiocaesium activity concentrations both in underground and aboveground parts of mycorrhizal sunflowers were nearly 10 fold greater when compared to those of nonmycorrhizal plants (see Fig. 1D). At the same time, roots of both mycorrhizal and nonmycorrhizal sunflowers had about 50 % higher ^{134}Cs activity concentrations when compared to plant shoots.

The presence of ^{134}Cs didn't have appreciable impact on *H. annuus* growth parameters, and the most distinct was the mycorrhiza influence. Thus, the mycorrhizal *H. annuus* grown on radioactive and clean substrata produced correspondingly 12 and 11 % longer shoots as compared to those of nonmycorrhizal plants. The shoots dry weight of mycorrhizal *H. annuus* grown both on radioactively contaminated and non-polluted soil exceeded substantially (about 70 and 80 % respectively) shoots dry weight of nonmycorrhizal ones. The degrees of ^{134}Cs translocation from roots to shoots of mycorrhizal and nonmycorrhizal *H. annuus* weren't differed considerably due to similar root/shoot ratios of the radionuclide activity concentration (1.54 ± 0.10 and 1.47 ± 0.21 correspondingly).




Fig. 1. ^{134}Cs activity concentration ($\text{Bq}\cdot\text{kg}^{-1}$) in roots and shoots of *Plantago lanceolata* (A), *Medicago truncatula* (B), *Lolium perenne* (C) and *Helianthus annuus* (D) mycorrhizal or not with *Glomus intraradices* and grown on substrata spiked with ^{134}Cs ($77\,000 \text{ Bq}\cdot\text{kg}^{-1}$). The results are presented as mean \pm standard deviation. The different letters above bars mean statistically significant differences ($p < 0.05$).

Part 2. Functioning of plant photosynthetic apparatus.

The most of photosynthesis biophysical parameters both in mycorrhizal and nonmycorrhizal plants cultivated on substrata with ^{134}Cs weren't varied considerably as compared to those of control ones (Fig. 2A-2D). The exception was observed only in case of *M. truncatula* grown on radioactively contaminated substrata which had considerably lower efficiency of trapped exciton movement into electron transport chain ($\psi_{E_0} = \text{ET}_0/\text{TR}_0$) and maximum yield of electron transport ($\phi_{E_0} = \text{ET}_0/\text{ABS}$) when compared to those of control plants from clean substrata. Also, the total and absorption vitality indexes (PI_{abs} and PI_{total}) of these alfalfas were respectively 33.2 ± 12.9 and 42.0 ± 18.7 % lower than those of control plants (Fig. 2B).




Fig. 2. Biophysical parameters of photosynthesis of *Plantago lanceolata* (A), *Medicago truncatula* (B), *Lolium perenne* (C) and *Helianthus annuus* (D): nonmycorrhizal (control) plants grown on clean soil (-Cs, -M) and plants mycorrhizal with *Glomus intraradices* and cultivated on substrata spiked with ^{134}Cs (+Cs, +M). Values on plots are presented in relative units and normalised on those of the control plants. * - means statistically significant difference ($p < 0.05$).

The fungal colonization of plant species grown on radioactive substrata had positive impact on functioning of *H. annuus* photosynthetic apparatus (Fig. 2D). In this case, the AM inoculation of plants cultivated on radioactive soil improved vitality indexes of sunflowers. Thus, PI_{abs} of mycorrhizal *H. annuus* was correspondingly 41.4 ± 11.3 and 36.3 ± 12.5 % higher than those of nonmycorrhizal plants grown

on soil with ^{134}Cs and control plants. In turn, PI_{total} of mycorrhizal sunflowers exceeded considerably those of nonmycorrhizal plants cultivated on radioactive substrata as well as control plants (40.0 ± 9.7 and 44.8 ± 11.9 % respectively, Fig. 2D).

Part 3. Arbuscular mycorrhizal colonization of plants.

Both treated with radiocaesium and control plant species were characterized with high mycorrhizal frequency (F%) that exceeded 90 %. The intraradical structures of the AM fungus were morphologically typical for Arum-type mycorrhizae. The intraradical hyphae of *G. intraradices* propagated between cortical root cells at long distances and formed lateral branches, which penetrated cells and produced arbuscules inside them (Fig. 3). The presence of numerous intercellular vesicles was characteristic for nearly 80 % of studied root fragments. The spores of *G. intraradices* that have thicker walls in comparison with vesicles were found only in several root fragments.

The most of AM colonization parameters of plants cultivated on substrata spiked with ^{134}Cs and non-polluted soil weren't differed significantly (Fig. 4A,C,D). Although, in case of *M. truncatula* the presence of radiocaesium resulted to considerable (about 30 %) decrease of mycorrhizal colonization intensity for all and individual mycorrhizal plant roots (M,% and m,% correspondingly) when compared to those of control plants (Fig. 4B).





Fig. 3. Arbuscules and vesicles of *Glomus intraradices* within roots of *Plantago lanceolata* (A) and *Medicago truncatula* (B) cultivated on substrata treated with ^{134}Cs ($77\,000 \text{ Bq}\cdot\text{kg}^{-1}$).






Fig. 4. Arbuscular mycorrhizal colonization parameters of *Plantago lanceolata* (A), *Medicago truncatula* (B), *Lolium perenne* (C) and *Helianthus annuus* (D) inoculated with *Glomus intraradices*: F% - frequency of mycorrhiza; M% - mycorrhizal colonization intensity for all roots; m% - mycorrhizal colonization intensity within individual mycorrhizal roots; A% - arbuscular richness for all roots; a% - arbuscular richness in root fragments where the arbuscules were present, medians. Plants were cultivated on clean substrata (-Cs) and substrata spiked with ^{134}Cs (+Cs). The different letters above bars mean statistically significant differences ($p < 0.05$).

The principal goals of the research were to compare the possible influence of mycorrhiza on various AM fungal symbionts cultivated on the same substrata under the impact of ^{134}Cs . Our results suggest that inoculation with AMF changed substantially the uptake of ^{134}Cs by studied plant species and influ-

enced the translocation of caesium isotopes within the plants. The arbuscular mycorrhiza resulted to considerable decrease of ^{134}Cs activity concentration in shoots of *P. lanceolata*, *M. truncatula* and *L. perenne* when compared to nonmycorrhizal ones. The most significant (about threefold) reduction of ^{134}Cs activity concentration was found in shoots of mycorrhizal *L. perenne*. This result contradicts to the data obtained by [9] who found that inoculation with arbuscular mycorrhiza significantly enhanced uptake of ^{137}Cs by ryegrass.

The exception in our study was *H. annuus* where the AM colonization led to nearly tenfold increase of ^{134}Cs activity concentration both in plant roots and shoots. The sunflower was previously shown to be an effective hyperaccumulator of ^{137}Cs and ^{60}Co [10], although the ability of this plant to form mycorrhiza has not been studied. In our case *H. annuus* revealed its ability of ^{134}Cs hyperaccumulation only in the presence of the mycorrhiza. Also, *H. annuus* was only plant species in our experiment whose shoot biomass was significantly affected by the impact of AM fungus. Such contradictory findings demonstrate that basic knowledge of Cs potential uptake mechanisms are needed to facilitate the design of countermeasures to reduce or enhance the transfer of radiocaesium into plants.

Conclusions. In summary, *M. truncatula* was suggested to be the most sensitive plant species relative to the radiocaesium impact. Due to considerable reduction of caesium in their shoots this plant species as well as *P. lanceolata* and *L. perenne* couldn't be applied in phytoremediation, but they may be potentially used in phytostabilization of the radioactively polluted ecosystems.

On the other hand, the use of *H. annuus* with its Cs hyperaccumulation properties conditioned by mycorrhiza for the phytoremediation is also questionable. In our study, the evaluated total activity of ^{134}Cs accumulated in biomass of sunflowers grown in one pot (two plants) during three months was 221 Bq. This is only 2.2 % from total radiocaesium activity in the pot (100 000 Bq). Extrapolating these data for a longer term and assuming the plant active growth period is about 6 months per year, we can roughly estimate that nearly two decades are needed to remove radiocaesium completely from the soil. This assumption doesn't take into consideration the natural factors, such as the radiocaesium migration, inhomogeneous distribution in soil and possible leaching of the radionuclide below the 30 - 40 cm (i.e. outside of root zone) as well as potential impact of another AM fungi and various soil microorganisms on the radionuclide uptake by plants.

The number of authors proposed using of AM fungi in phytoremediation strategies for radiocaesium contaminated areas to enhance radionuclide re-

moval by plant biomass [1, 11, 12]. Although, the effects of AM fungi on Cs accumulation could be applied in strategies to develop crops with smaller soil-to-plant transfer factors which accumulate less Cs [8, 12]. Such plant species may be potentially grown within areas with moderate radiocaesium contamination levels and further used in agricultural purposes, if the radionuclide content in their biomass doesn't exceed the prescribed permissible levels. Our results demonstrated the capacity of AM fungi to influence the acquisition and accumulation of caesium isotopes by plants by immobilizing, transporting and affecting the root-to-shoot translocation. Nevertheless the AM fungal ability to take part in phytoremediation strategies still remains questionable and needs for further researches.

Acknowledgements. This research project was supported by a Marie Curie Early Stage Research Training Fellowship of the European Community's Sixth framework Programme under contract number MEST-CT-2005-020387.

References

1. Entry J.A., Vance N.C., Hamilton M.A. et al. Phytoremediation of soil contaminated with low concentrations of radionuclides // *Water Air Soil Pollut.* - 1996. - Vol. 88. - P. 167 - 176.
2. Galanda D., Matel L., Strisovska J., Dulanska S. Mycoremediation: the study of transfer factor for plutonium and americium uptake from the ground// *J. Radioanal. Nucl. Chem.* - 2014. - Vol. 299. - P. 1411 - 1416.
3. Entry J.A., Astrud L.S., Reeves M. Accumulation of ^{137}Cs and ^{90}Sr from contaminated soil by three grass species inoculated with mycorrhizal fungi // *Environ. Pollut.* - 1997. - Vol. 104. - P. 449 - 457.
4. Dupré de Boulois H., Joner E.J., Leyval C. et al. Role and influence of mycorrhizal fungi on radiocaesium accumulation by plants // *J. Environ. Radioact.* - 2008. - Vol. 99. - P. 785 - 800.
5. Chen B.D., Jakobsen I., Roos P., Borgaard O.K., Zhu Y.G. Mycorrhiza and root hairs enhance acquisition of phosphorus and uranium from phosphate rock but mycorrhiza decreases root to shoot uranium transfer // *New Phytol.* - 2005 - Vol. 165. - P. 591 - 598.
6. Trouvelot A., Kough J.L., Gianinazzi-Pearson V. Mesure du taux de mycorrhization VA d'un système racinaire. Recherche de méthodes d'estimation ayant une signification

- fonctionnelle // Physiological and Genetical Aspects of Mycorrhizae // INRA Press Paris. – 1986. – P. 217 – 221.
7. Strasser R.J., Tsimilli-Michael M., Srivastava A. Analysis of the chlorophyll a fluorescence transient // Chlorophyll a fluorescence: a signature of photosynthesis. Advances in Photosynthesis and Respiration Series. Kluwer Academic Publishers. Rotterdam. – 2004. – Vol. 19. – P. 321 – 362.
 8. Gyuricza V., Declerck S., Dupré de Boulois H. Arbuscular mycorrhizal fungi decrease radiocesium accumulation in *Medicago truncatula* // J Environ Radioact. – 2010. – Vol. 101. – P. 591 – 596.
 9. Rosen, K., Zhong W.L., Martensson A. Arbuscular mycorrhizal fungi mediated uptake of ¹³⁷Cs in leek and ryegrass // Sci. Total Environ. – 2005. – Vol. 338, No. 3. – P. 283 – 290.
 10. Hornik M., Pipiska M., Vrtoch L. et al. Bioaccumulation of Cs-137 and Co-60 by *Helianthus annuus* // Nukleonika. – 2005. – Vol. 50. – P. 49 – 52.
 11. Dubchak S., Ogar A., Mietelski J.W., Turnau K. Influence of silver and titanium nanoparticles on arbuscular mycorrhiza colonization and accumulation of radiocesium in *Helianthus annuus* // Span J Agric Res. – 2010. – Vol. 8. – P. 103 – 108.
 12. Willey N. Amelioration of soils contaminated with radionuclides: Exploiting biodiversity to minimize or maximise soil to plant transfer // Radioprotection. – 2005. – Vol. 40. – P. 819 – 824.

ЕКОЛОГІЯ І ВИРОБНИЦТВО

УДК 504.064; 614.835.3

ТЕХНОГЕННІ РИЗИКИ ЗАБРУДНЕННЯ ДОВКІЛЛЯ ПІД ЧАС РЕМОНТНИХ РОБІТ РЕЗЕРВУАРІВ ІЗ НАФТОПРОДУКТАМИ

Липовий В. О., Удянський М. М.
Національний університет цивільного захисту України,
вул. Чернишевська, 94, 61023, м. Харків
lipovoy_vladimir@mail.ru

Наведено інформацію щодо поліпшення стану забезпечення екологічної безпеки навколо небезпечних об'єктів з наявністю технологічних процесів, пов'язаних з експлуатуванням та проведенням ремонтних робіт резервуарів із нафтопродуктами шляхом управління техногенними ризиками з урахуванням впливу чинників на їх значення. *Ключові слова:* техногенний ризик, забруднення довкілля, резервуар з нафтопродуктами, пробіт-функція, нафтозалишки.

Техногенные риски загрязнения окружающей среды при проведении ремонтных работ резервуаров с нефтепродуктами. Липовой В.А., Удянский Н.Н. Приведена информация об улучшении состояния обеспечения экологической безопасности вокруг опасных объектов с наличием технологических процессов, связанных с эксплуатацией и проведением ремонтных работ резервуаров с нефтепродуктами, путем управления техногенными рисками с учетом влияния факторов на их значение. *Ключевые слова:* техногенный риск, загрязнение окружающей среды, резервуар с нефтепродуктами, пробит-функция, нефтеостаток.

Man-caused environmental pollution during repair tanks with oil. Lipovoy V.O., Udyansky N.N. The data on the improvement of environmental safety around dangerous objects to the presence of processes associated with the operation and maintenance work tanks with oil by controlling technological risks, taking into account certain factors influence their value. *Keywords:* technological hazards, pollution, oil reservoir, the probit function, the residue oil.

Аварійні викиди та витіки шкідливих речовин внаслідок проведення регламентних та ремонтних робіт з очищення внутрішніх технологічних поверхонь резервуарів із нафтопродуктами можуть призвести до локального та катастрофічного рівня за-
вдань шкоди довкіллю та життєдіяльності людей.
Статистика свідчить, що понад 20% усіх пожеж на резервуарах зберігання нафтопродуктів відбувається через порушення вимог пожежної безпеки при проведенні ремонтних